arXiv:1603.02864v1 [math.SG] 9 Mar 2016

THE AUTONOMOUS NORM ON $\mathrm{Ham}(\mathbf{R}^{2n})$ IS BOUNDED

MICHAEL BRANDENBURSKY AND JAREK KĘDRA

Abstract. We prove that the autonomous norm on the group of compactly supported Hamiltonian diffeomorphisms of the standard $\mathbf{R}^{2n}$ is bounded.

Let (M,ω) be a symplectic manifold and let $\mathrm{Ham}(M,\omega)$ be the group of compactly supported Hamiltonian diffeomorphisms of (M,ω). Recall that a Hamiltonian diffeomorphism f is a time-one map of the flow generated by the vector field X_{F_t} defined by $\omega(X_{F_t},-)=dF_t$. Here $F\colon M\times S^1\to\mathbf{R}$ is a smooth compactly supported function and $F(x,t)=F_t(x)$ (see [9, Section 5.1] for details). The function F is called a Hamiltonian of f. If F does not depend on the second variable or is time independent then f is called autonomous. It is known that every Hamiltonian diffeomorphism is a product of autonomous ones [3]. The autonomous norm on $\mathrm{Ham}(M,\omega)$ is defined by:

$$\|f\|=\min\{k\in\mathbf{N}\mid f=a_1\cdots a_n,\ \text{where } a_i \text{ is autonomous}\}.$$

It is a conjugation invariant norm and is known to be unbounded on the group of compactly supported Hamiltonian diffeomorphisms of an oriented surface of finite area [2, 4, 3, 6].

This paper is concerned with the group $\mathrm{Ham}(\mathbf{R}^{2n})$ of compactly supported Hamiltonian diffeomorphism of the Euclidean space equipped with the standard symplectic form. We prove the following result.

Theorem. *The diameter of the autonomous norm on* $\mathrm{Ham}(\mathbf{R}^{2n})$ *is bounded above by* 3.

Proof. Let $f\in\mathrm{Ham}(\mathbf{R}^{2n})$. Let $f=a_1\cdots a_m$, where $a_i\in\mathrm{Ham}(\mathbf{R}^{2n})$ are autonomous diffeomorphisms with compactly supported Hamiltonian functions $F_i\colon\mathbf{R}^{2n}\to\mathbf{R}$. Let $B(r)\subset\mathbf{R}^{2n}$ be an Euclidean ball of radius $r>0$, centered at the origin and containing the union of the supports of the functions F_i.

Lemma. *There exists an autonomous diffeomorphism* $h\in\mathrm{Ham}(\mathbf{R}^{2n})$ *displacing the ball* $B(r)$ m *times.*

The statement of the lemma means that $h^i(B(r)) \cap h^j(B(r)) = \emptyset$ for $0 \leq i \neq j \leq m$. It follows from [5, Lemma 2.6] that there exists $g \in \mathrm{Ham}\,(\mathbf{R}^{2n})$ such that the following equality holds:

$$f = a_1 \cdots a_m = [h, g] a_1^h a_2^{h^2} \cdots a_m^{h^m},$$

where $a_i^{h^i} = h^i a_i h^{-i}$ and h is the diffeomorphism from the Lemma. Observe that, since the supports of $F_i \circ h^i$ are pairwise disjoint for $i \in \{1, \ldots, m\}$, we obtain that the composition $a_1^h a_2^{h^2} \cdots a_m^{h^m}$ is autonomous with the Hamiltonian function equal to

$$F_1 \circ h + F_2 \circ h^2 + \cdots + F_m \circ h^m.$$

Since the commutator $[h, g] = h \cdot h^g$ is a product of two autonomous diffeomorphisms we obtain that f is a product of three autonomous diffeomorphisms. □

Proof of the Lemma. Let $H_1 \colon \mathbf{R} \to \mathbf{R}$ be a smooth function satisfying the following conditions:

(1) $H_1(y) = 0$ for $|y| > r + 1$

(2) $H_1'(y) = r$ for $|y| \leq r$.

Let $H(x_1, y_1, \ldots, x_n, y_n) = H_1(y_1)$. We have that $dH = r dx_1$ and that the induced Hamiltonian vector field X is equal to $r\frac{\partial}{\partial x_1}$. Thus the induced Hamiltonian diffeomorphism displaces the ball $B(r)$ as many times as we like. Taking an appropriate cut off function we obtain the required compactly supported diffeomorphism h. □

Remarks. If f in Theorem is contained in the kernel of the Calabi homomorphism (see Section 8.B of [1] for a definition) then the same argument shows that it is a product of up to three autonomous diffeomorphisms with trivial Calabi invariant.

It is known that the Hofer norm on $\mathrm{Ham}\,(\mathbf{R}^{2n})$ is unbounded and stably bounded [7]. The kernel of the Calabi homomorphism does not admit nontrivial quasimorphisms, however, it is stably unbounded [8].

It is not difficult to see that the diameter of the autonomous norm on $\mathrm{Ham}\,(\mathbf{R}^{2n})$ is at least 2. To the best of our knowledge it is an open question whether there exists a Hamiltonian diffeomorphism of $\mathbf{R}^{2n}$ of autonomous norm equal to 3.

Acknowledgements. We thank the Center for Advanced Studies in Mathematics at Ben Gurion University for supporting the visit of the second author at BGU.

References

[1] Vladimir I. Arnold and Boris A. Khesin. *Topological methods in hydrodynamics*, volume 125 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1998.

[2] Michael Brandenbursky and Jarek Kędra. On the autonomous metric on the group of area-preserving diffeomorphisms of the 2-disc. *Algebr. Geom. Topol.*, 13(2):795–816, 2013.

[3] Michael Brandenbursky, Jarek Kedra, and Egor Shelukhin. On the autonomous norm on the group of Hamiltonian diffeomorphisms of the torus. Available at http://arxiv.org/abs/1602.03287v2.

[4] Michael Brandenbursky and Egor Shelukhin. On the L^p-geometry of autonomous Hamiltonian diffeomorphisms of surfaces. *Math. Res. Lett. to appear*. Available at http://arxiv.org/abs/1405.7931v2.

[5] Dmitri Burago, Sergei Ivanov, and Leonid Polterovich. Conjugation-invariant norms on groups of geometric origin. In *Groups of diffeomorphisms*, volume 52 of *Adv. Stud. Pure Math.*, pages 221–250. Math. Soc. Japan, Tokyo, 2008.

[6] Jean-Marc Gambaudo and Étienne Ghys. Commutators and diffeomorphisms of surfaces. *Ergodic Theory Dynam. Systems*, 24(5):1591–1617, 2004.

[7] Helmut Hofer and Eduard Zehnder. *Symplectic invariants and Hamiltonian dynamics*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Verlag, Basel, 1994.

[8] Morimichi Kawasaki. Relative quasimorphisms and stably unbounded norms on the group of symplectomorphisms of the euclidean spaces. *Journal of Symplectic Geometry*, 2014.

[9] Leonid Polterovich. *The geometry of the group of symplectic diffeomorphisms*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2001.

Ben Gurion University, Israel

E-mail address: brandens@math.bgu.ac.il

University of Aberdeen and University of Szczecin

E-mail address: kedra@abdn.ac.uk